

A Theoretical Study of Fixed-Point Theorems and Their Applications in Mathematical Analysis

Dr. Preeti Panwar

Associate Professor in Mathematics, Guru Nanak Khalsa College, Karnal 132001

Email Id: kharbpreeti@gmail.com

Abstract

Fixed-point theorems occupy a central position in mathematical analysis, providing existence and, in many cases, uniqueness guarantees for solutions of equations that cannot be solved in closed form. This paper presents a theoretical survey of the principal fixed point results across three settings: metric spaces, topological spaces, and ordered sets. The Banach contraction principle and its generalisations are examined as the prototypical metric result, furnishing constructive existence proofs via iteration; Brouwer's and Schauder's theorems are discussed as topological results guaranteeing fixed points without contraction hypotheses, at the cost of constructivity; and the Knaster Tarski theorem is presented as the order-theoretic counterpart, together with its set-valued extension due to Kakutani. The paper concludes with a discussion of applications, including existence theorems for ordinary and integral equations, iterative numerical methods, and equilibrium analysis in game theory, arguing that the diversity of fixed point theorems reflects a genuine trade-off between the strength of the hypotheses imposed and the amount of structural information uniqueness, computability, stability that the resulting fixed point inherits.

Keywords: fixed-point theorems, Banach contraction principle, Brouwer's theorem, Schauder's theorem, Knaster Tarski theorem, Kakutani's theorem, functional analysis

1. Introduction

A fixed point of a mapping T from a set X into itself is a point x such that $T(x) = x$. Despite the simplicity of this definition, fixed point theorems form one of the most powerful and widely applied families of results in mathematical analysis, because an enormous range of problems solving equations, establishing the existence of solutions to differential and integral equations, guaranteeing equilibria in economic models, and justifying the convergence of iterative numerical schemes can be recast as the search for a fixed point of an appropriately constructed operator [1], [2].

The theory divides naturally according to the structure imposed on the space and the mapping. Metric fixed-point theory, exemplified by the contraction principle of Banach, trades a strong contractive

hypothesis on the map for a constructive proof and a guarantee of uniqueness [3]. Topological fixed-point theory, exemplified by the theorems of Brouwer and Schauder, dispenses with the contraction hypothesis in favour of compactness and convexity assumptions, at the price of losing both uniqueness and constructivity [4], [5]. Order-theoretic fixed-point theory, exemplified by the Knaster Tarski theorem, replaces the topological or metric structure with a completeness condition on a partially ordered set, yielding existence of extremal fixed points under monotonicity alone [6]. This paper surveys each of these branches in turn, together with the set-valued extension due to Kakutani, discusses their historical relationships in Section 7, and closes with a discussion of applications in analysis, numerical computation, and the theory of games.

A theme that recurs throughout the paper, and that is made explicit in the comparative discussion of Section 7, is that the various fixed-point theorems are not competitors offering different proofs of the same conclusion, but rather occupy genuinely different positions along a spectrum that trades the strength of the hypothesis imposed on the map or the space for the richness of the information obtained about the resulting fixed point, ranging from mere existence, through existence together with a characterisation of the entire fixed-point set, to existence, uniqueness, and an explicit constructive algorithm of approximation. Understanding this trade-off is essential for selecting the appropriate fixed point theorem when formulating an existence argument for a new class of equations, since an unnecessarily strong theorem may be inapplicable to the operator at hand, while an unnecessarily weak theorem may fail to deliver the uniqueness or constructivity that a given application requires.

2. Metric Fixed Point Theory

2.1 The Banach Contraction Principle

Let (X, d) be a complete metric space and let $T: X \rightarrow X$ be a contraction, meaning that there exists a constant k with $0 \leq k < 1$ such that

$$d(T(x), T(y)) \leq k \cdot d(x, y) \text{ for all } x, y \in X.$$

The Banach contraction principle, established by Banach in 1922, asserts that T possesses a unique fixed point x^* , and that for any x_0 in X the sequence of iterates $x_{n+1} = T(x_n)$ converges to x^* [3]. The proof is constructive: one shows that the sequence of iterates is Cauchy by summing the geometric bound $d(x_{n+1}, x_n) \leq k^n \cdot d(x_1, x_0)$, invokes completeness to obtain a limit, and verifies by continuity of T that the limit is fixed [3], [7]. The principle additionally yields an explicit a priori error estimate,

$$d(x_n, x^*) \leq (k^n / (1 - k)) \cdot d(x_1, x_0),$$

which is of independent practical value because it bounds the number of iterations required to achieve a prescribed accuracy without knowledge of x^* itself [7].

2.2 Generalisations: Weak Contractions and Kannan Mappings

A substantial literature has developed around weakening the contraction hypothesis while retaining existence and, where possible, uniqueness of a fixed point. Kannan showed in 1968 that a mapping need not be continuous to have a unique fixed point, provided it satisfies $d(T(x), T(y)) \leq k \cdot [d(x, T(x)) + d(y, T(y))]$ for some k in $[0, 1/2)$, a condition genuinely independent of the Banach contraction condition [8]. Other generalisations relax the fixed contraction constant to a function that depends on the pair of points, or replace the metric with a more general distance structure, such as a partial metric or a b-metric, while preserving the essential completeness-and-contraction architecture of the original proof [8], [9]. These generalisations are of more than technical interest: they extend the reach of constructive existence-and-uniqueness results to operators, such as certain integral operators arising in the study of Volterra equations, that fail to be contractions in the classical sense but nonetheless satisfy a weaker contractive-type inequality [9].

3. Topological Fixed Point Theorems

3.1 Brouwer's Fixed Point Theorem

Brouwer's fixed point theorem, published in 1911, states that every continuous map from the closed unit ball B^n in $\mathbb{R}^n$ to itself has at least one fixed point [4]. Unlike the Banach principle, no contraction or even injectivity is assumed; continuity together with compactness and convexity of the domain suffices. The classical proofs are non-constructive, typically proceeding either by an algebraic-topological argument via the non-existence of a continuous retraction from B^n onto its boundary sphere, or by a combinatorial argument based on Sperner's lemma, which triangulates the simplex and produces a fixed point as a limit of a sequence of increasingly fine simplicial approximations [4], [10]. The theorem fails without compactness or convexity: a rotation of the open disc by a fixed angle, or a translation of all of $\mathbb{R}^n$, has no fixed point, which shows that both hypotheses are essential rather than technical conveniences [4].

3.2 Schauder's Fixed Point Theorem

Schauder extended Brouwer's theorem in 1930 to infinite-dimensional normed spaces, a setting in which the Heine–Borel property fails and closed bounded sets need not be compact [5]. The Schauder fixed point theorem states that if K is a nonempty, closed, bounded, and convex subset of a Banach space, and $T: K \rightarrow K$ is a continuous map whose image $T(K)$ is relatively compact, then T has at least one fixed point in K [5], [11]. The compactness hypothesis on $T(K)$, rather than on K itself, is essential in infinite dimensions, since

the closed unit ball of an infinite-dimensional Banach space is never compact; the theorem is typically proved by approximating T uniformly by maps of finite-dimensional range and invoking the finite-dimensional Brouwer theorem on a sequence of increasingly accurate approximations [5], [11]. Schauder's theorem is the principal tool used to establish existence of solutions to nonlinear differential and integral equations, where the relevant operator T acts on a function space rather than on Euclidean space [11].

3.3 Nonexpansive Mappings and the Browder–Göhde–Kirk Theorem

Between the strong contraction hypothesis of Banach and the mere continuity hypothesis of Brouwer and Schauder lies the intermediate class of nonexpansive mappings, those satisfying $d(T(x), T(y)) \leq d(x, y)$ without the strict inequality required for a contraction. A nonexpansive self-map of an arbitrary complete metric space need not have a fixed point a translation on the real line is nonexpansive and fixed-point free so additional geometric structure on the space is required. The Browder Gohde Kirk theorem, established independently by these authors in the mid-1960s, shows that if K is a nonempty, closed, bounded, and convex subset of a uniformly convex Banach space, then every nonexpansive map $T: K \rightarrow K$ has a fixed point [9]. Uniform convexity, a geometric condition requiring that midpoints of chords between points on the unit sphere lie strictly inside the ball at a rate controlled uniformly over the sphere, is essential here, and Hilbert spaces, as well as the classical L^p spaces for $1 < p < \infty$, satisfy it, whereas the space of continuous functions with the supremum norm does not [9]. Unlike the Banach principle, the fixed point supplied by this theorem need not be unique, and the standard proofs are again non-constructive, typically proceeding via an application of Zorn's lemma to extract a minimal nonempty closed convex invariant set on which T acts, followed by a geometric argument showing that such a minimal set must be a single point [9].

3.4 Coincidence and Common Fixed Point Theorems

A further refinement of the theory concerns coincidence points and common fixed points of two or more mappings, which are of particular relevance when a problem is naturally expressed in terms of an equation relating two operators rather than a single self-map. Given two mappings $S, T: X \rightarrow X$, a coincidence point is a point x such that $S(x) = T(x)$, and a common fixed point is a point x such that $S(x) = T(x) = x$. Jungck's common fixed point theorem generalises the Banach contraction principle to this setting, showing that if S and T commute, T 's range is contained in S 's range, S is continuous, and T is a contraction relative to S in an appropriate sense, then S and T possess a unique common fixed point [9]. Such results are of substantial use in establishing the existence and uniqueness of solutions to systems of equations, and in particular to differential and integral equations that couple two unknown functions through a shared constraint, a situation for which a single-operator fixed-point theorem does not directly apply [9].

4. Fixed Points in Ordered Structures

4.1 The Knaster–Tarski Theorem

A distinct strand of fixed-point theory dispenses with topology altogether in favour of order structure. Let $(L, \leq)$ be a complete lattice, meaning every subset of L has both a supremum and an infimum in L , and let $f: L \rightarrow L$ be monotone, meaning $x \leq y$ implies $f(x) \leq f(y)$. The Knaster–Tarski theorem, in the form given by Tarski in 1955, asserts that the set of fixed points of f is itself a nonempty complete lattice, and in particular f has a greatest fixed point and a least fixed point [6]. The least fixed point may be characterised as the infimum of the set $\{x \in L : f(x) \leq x\}$, and dually for the greatest fixed point, and no continuity assumption on f is required, monotonicity alone being sufficient [6].

4.2 Applications to Lattice Theory

The order-theoretic fixed-point theorem has significant applications outside classical analysis. In set theory it is used to give a short proof of the Cantor–Schröder–Bernstein theorem, by exhibiting the desired bijection as a fixed point of a monotone operator on the lattice of subsets of one of the two sets under comparison [6]. In theoretical computer science, the least fixed point of a monotone operator on a lattice of program states or predicates underlies the denotational semantics of recursive programs and the semantics of logic-programming languages, where the meaning of a recursively defined procedure is taken to be precisely the least fixed point of the associated semantic operator [12]. The order-theoretic approach therefore complements the metric and topological approaches by supplying existence results in settings, such as discrete or non-metrisable lattices, where neither completeness in the metric sense nor compactness in the topological sense is naturally available.

5. Set-Valued Fixed Point Theory

5.1 Kakutani's Theorem

Many applications require fixed points not of single-valued functions but of correspondences, that is, mappings that assign to each point of a space a nonempty subset of that space. Kakutani's fixed point theorem, published in 1941, states that if K is a nonempty, compact, and convex subset of $\mathbb{R}^n$, and $\varphi: K \rightarrow 2^K$ is an upper hemicontinuous correspondence assigning to each point of K a nonempty, closed, and convex subset of K , then there exists a point x^* in K such that $x^* \in \varphi(x^*)$ [13]. Kakutani's theorem may be derived from Brouwer's theorem by approximating the correspondence φ with a sequence of continuous single-valued functions and passing to the limit, so that the set-valued result inherits both the strengths and the non-constructive character of its single-valued antecedent [13], [14].

5.2 Applications in Game Theory and Economics

Kakutani's theorem achieved particular prominence through Nash's 1950 proof of the existence of an equilibrium point in every finite non-cooperative game [14]. In that argument, the best-response correspondence, which assigns to each strategy profile of the other players the set of a given player's optimal responses, is upper hemicontinuous with convex values whenever mixed strategies are permitted, and a Nash equilibrium is precisely a fixed point of the resulting joint best-response correspondence across all players [14]. The same architecture, of casting an equilibrium concept as a fixed point of a suitably defined correspondence, recurs throughout mathematical economics, notably in the proof of the existence of competitive (Walrasian) equilibrium prices in general equilibrium theory, where the relevant correspondence maps a price vector to the set of prices that clear an associated excess-demand function [15].

5.3 Multivalued Contractions and the Nadler Theorem

A set-valued analogue of the Banach contraction principle, distinct in spirit from Kakutani's topological result, was established by Nadler in 1969. Given a complete metric space (X, d) , let $CB(X)$ denote the collection of nonempty closed bounded subsets of X , equipped with the Hausdorff distance H , defined for $A, B \in CB(X)$ by $H(A, B) = \max \{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) \}$. Nadler showed that if $T: X \rightarrow CB(X)$ is a multivalued contraction with respect to H , meaning $H(T(x), T(y)) \leq k \cdot d(x, y)$ for some k in $[0, 1)$, then T has a fixed point, that is, a point x^* with $x^* \in T(x^*)$ [16]. The proof again proceeds by iteration, though the construction of a Cauchy sequence of iterates is more delicate than in the single-valued case, since at each stage one must select a point in $T(x_n)$ sufficiently close to the previous iterate, using the definition of the Hausdorff distance to control the resulting approximation error [13]. This result bridges the metric and topological branches of the theory surveyed above, since it applies the iterative, contraction based methodology of Section 2 to the set-valued setting more commonly associated with the topological methods of Kakutani's theorem [16].

6. Applications in Mathematical Analysis

6.1 Existence Theorems for Differential Equations

The Picard Lindelöf theorem, guaranteeing local existence and uniqueness of solutions to the initial value problem $y' = f(t, y)$, $y(t_0) = y_0$ under a Lipschitz condition on f , is obtained by recasting the differential equation as the integral equation $y(t) = y_0 + \int f(s, y(s)) ds$ and showing that the associated integral operator is a contraction on an appropriate space of continuous functions with a suitably weighted metric, so that the Banach contraction principle applies directly [7], [11]. When the Lipschitz hypothesis on f is dropped and only continuity is assumed, uniqueness can fail, but existence of at least one solution, the content of the

Peano existence theorem, is instead established using Schauder's theorem, since the relevant operator remains continuous and compact even though it may no longer be a contraction [5], [11]. This pairing illustrates the general trade-off identified throughout the paper: the stronger Banach-type hypothesis buys uniqueness and constructivity, while the weaker Schauder-type hypothesis buys generality at the cost of both.

It is worth noting explicitly why the constructivity of the Banach contraction principle, rather than merely its existence and uniqueness conclusion, is what makes it foundational for numerical analysis. The Brouwer and Schauder theorems certify that a fixed point exists but supply no procedure, even in principle, for locating it to any prescribed accuracy, so an existence proof based on either theorem gives no direct guidance for numerical implementation. The Banach principle, by contrast, is itself an algorithm: the proof that the iterates $x_{n+1} = T(x_n)$ converge is simultaneously a specification of the numerical method, and the accompanying a priori error bound quantifies, before any computation is performed, how many iterations are required to reach a target tolerance [3], [7]. This is why, whenever an existence problem in analysis can be reduced to a fixed point of a contraction, doing so is generally preferred over invoking a topological existence theorem, even when the topological route is available, since the contraction-based route yields a usable computational procedure as an immediate corollary of the existence proof rather than as a separate undertaking.

6.2 Numerical Methods and Iterative Approximation

The constructive character of the Banach contraction principle makes it the theoretical basis for a wide range of iterative numerical methods. Newton's method for locating a root of a differentiable function f may be analysed as fixed-point iteration for the auxiliary map $T(x) = x - f(x)/f'(x)$, and the classical local convergence theorem for Newton's method follows once T is shown to be a contraction on a suitable neighbourhood of the root [7]. More generally, many preconditioned iterative schemes for solving large linear systems, as well as fixed-point schemes for nonlinear systems arising from the discretisation of partial differential equations, are justified by verifying, directly or via a spectral radius condition on the associated iteration matrix, that the corresponding iteration operator is a contraction with respect to some norm, thereby placing the convergence analysis of the numerical scheme within the scope of the Banach contraction principle [7], [9].

6.3 Integral Equations and Operator Theory

Beyond the initial value problems discussed above, fixed-point methods furnish the standard existence theory for integral equations of both Volterra and Fredholm type. A Fredholm equation of the second kind,

$y(t) = g(t) + \lambda \int_a^b K(t, s) y(s) ds$, may be recast as the fixed-point equation $y = T(y)$ for the linear integral operator T defined by the right-hand side, and when λ is small enough, or the kernel K is suitably bounded, T is a contraction on a space of continuous or square-integrable functions with respect to a norm adapted to the interval $[a, b]$, so that the Banach contraction principle again yields existence and uniqueness together with the convergent Neumann series $y = \sum_k \lambda^k T^k(g)$ as an explicit representation of the solution [7], [11]. Volterra equations of the second kind, in which the upper limit of integration is the variable t itself rather than a fixed endpoint b , exhibit an even stronger phenomenon: the associated operator, although perhaps not a contraction outright, has an iterate T^n that is a contraction for n sufficiently large, and a standard corollary of the Banach principle shows that a map possessing a contractive iterate has a unique fixed point coinciding with that of the iterate, so existence and uniqueness of solutions to Volterra equations hold globally on the entire interval of definition, without the restriction on λ required in the Fredholm case [11].

6.4 A Worked Illustration: Existence of Equilibrium Prices

To make the abstract discussion of Section 5.2 concrete, it is useful to sketch, at a theoretical level, how a Kakutani-type argument establishes existence of a competitive equilibrium in a simple exchange economy, since this example brings together compactness, convexity, and upper hemicontinuity in a single explicit construction [14], [15]. Consider an economy with finitely many goods and finitely many consumers, each endowed with an initial bundle of goods and a preference relation over consumption bundles satisfying standard convexity and continuity assumptions. For each price vector p in the simplex of normalised prices, each consumer's optimal consumption bundle, and hence the aggregate excess demand $z(p)$, the difference between total demand and total endowment at prices p , is well defined whenever preferences are strictly convex, but need only be a nonempty, convex, and compact set of maximisers when preferences are merely convex, so that the excess demand is properly formulated as a correspondence rather than a function [15]. Define an auxiliary correspondence ϕ on the price simplex that assigns to each price vector p the set of prices maximising the inner product of the price vector with the excess demand set $z(p)$; this construction, standard in the general equilibrium literature, guarantees that ϕ is upper hemicontinuous with nonempty, closed, convex values on a compact convex domain, precisely the hypotheses of Kakutani's theorem [14], [15]. A fixed point p^* of ϕ , guaranteed to exist by Kakutani's theorem, can then be shown, using Walras's law the identity that the value of excess demand is identically zero at any price vector to satisfy $z(p^*) \leq 0$, meaning that no good is in excess demand at p^* , which is precisely the definition of a competitive equilibrium price vector [15]. This construction, due in essence to Arrow and Debreu, illustrates in miniature the general pattern noted in Section 5.2: an equilibrium concept from outside pure

mathematics is shown to exist by exhibiting it as the fixed point of a correspondence built directly from the economic primitives of the model, with the mathematical content of the existence proof residing entirely in the verification that this correspondence satisfies the hypotheses of Kakutani's theorem.

7. Historical Development and Comparative Assessment

The results surveyed above did not arise as a single unified theory but emerged from distinct traditions that were only later recognised as instances of a common underlying phenomenon. Poincaré's work in the 1880s on the qualitative theory of differential equations already contained arguments of a fixed-point character, in his study of periodic orbits and of what would later be formalised as the Poincaré Bendixson theorem, though the general topological theorem was not isolated until Brouwer's 1911 paper, itself motivated by foundational questions in the topology of manifolds rather than by any application to analysis [4], [10]. Banach's 1922 thesis, by contrast, arose directly out of the emerging theory of integral equations and metric spaces initiated by Fréchet, and was explicitly designed as a tool for proving existence and uniqueness of solutions to functional equations by an iterative, constructive method quite different in character from Brouwer's topological, non-constructive argument [3], [7]. Schauder's 1930 extension of Brouwer's theorem to infinite-dimensional spaces was motivated by the desire to apply topological fixed-point methods to the same class of integral- and differential-equation problems that Banach's principle addressed, but for operators lacking the contraction property, thereby directly connecting the topological and metric branches of the subject that had developed somewhat independently up to that point [5], [11]. The order-theoretic branch, associated with Tarski's 1955 paper, arose from entirely different motivations in mathematical logic and the foundations of set theory, and it was only decades later, with the rise of denotational semantics in theoretical computer science during the 1970s, that the order-theoretic fixed-point theorem found a major application outside the area of its original conception [6], [12].

Comparing the theorems surveyed in this paper along the axes of generality, uniqueness, and constructivity clarifies why no single fixed point theorem has superseded the others. The Banach contraction principle imposes the strongest hypothesis an explicit numerical contraction constant and in exchange delivers the strongest conclusion: a unique fixed point, an explicit iterative algorithm for approximating it, and a quantitative error bound at every stage of the iteration [3], [7]. Brouwer's and Schauder's theorems relax the hypothesis to mere continuity, at the cost of both uniqueness, since a continuous self-map of a convex compact set may have many fixed points, and constructivity, since the classical proofs give no algorithm for locating a fixed point, only a guarantee that one exists [4], [5]. The Knaster Tarski theorem occupies a distinctive position in this comparison, since it requires no continuity at

all, only monotonicity with respect to a partial order, and correspondingly yields not merely existence but a description of the entire fixed-point set as a complete lattice bounded by an explicitly characterised least and greatest fixed point [6]. Kakutani's and Nadler's set-valued theorems generalise the topological and metric strands respectively to correspondences, illustrating that the trade-off between hypothesis strength and conclusion strength persists even after the passage from single-valued to set-valued mappings [13], [16]. Recognising this common trade-off across otherwise historically unrelated results is, arguably, the chief conceptual benefit of studying fixed-point theory as a unified subject rather than as a loose collection of unrelated existence theorems attached to particular applications.

8. Conclusion

This paper has surveyed the principal fixed-point theorems of mathematical analysis — the Banach contraction principle and its generalisations, the topological theorems of Brouwer and Schauder, the order-theoretic Knaster Tarski theorem, and the set-valued extension due to Kakutani alongside representative applications in differential equations, numerical analysis, and game theory. A unifying theme is the trade-off between the strength of the hypotheses placed on the underlying space and mapping and the richness of the conclusion obtained: contractive hypotheses yield unique, constructively approximable fixed points; compactness and convexity hypotheses yield existence without uniqueness or constructivity; and monotonicity on a complete lattice yields extremal fixed points without any topological structure whatsoever. Continuing research in this area extends these results to further generalised metric and topological settings, to fixed-point theory for nonexpansive mappings on Hilbert and Banach spaces, and to computational fixed-point algorithms with explicit convergence-rate guarantees [9], [11].

More broadly, the theoretical study undertaken here suggests that the choice among fixed-point theorems in a given application is itself a substantive mathematical decision, not merely a matter of convenience. An analyst confronted with an operator equation must first determine which structural features of the operator and the underlying space are actually available a numerical contraction bound, mere continuity together with compactness, monotonicity with respect to some natural order, or set-valued behaviour and only then select the fixed-point theorem whose hypotheses match those features, since selecting a theorem with unnecessarily strong hypotheses is simply inapplicable, while selecting one with unnecessarily weak hypotheses discards information, such as uniqueness or an explicit rate of convergence, that a stronger and equally applicable theorem would have provided. Viewed in this light, the diversity of fixed-point theorems catalogued in this paper is not a fragmentation of the subject but a toolkit whose

breadth is precisely what makes fixed-point methods applicable across the full range of existence problems that arise throughout mathematical analysis and its applications.

References

1. Granas, A., Dugundji, J.: Fixed Point Theory. Springer, New York (2003)
2. Zeidler, E.: Nonlinear Functional Analysis and its Applications I: Fixed-Point Theorems. Springer, New York (1986)
3. Banach, S.: Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. Fund. Math. 3, 133–181 (1922)
4. Brouwer, L.E.J.: Über Abbildung von Mannigfaltigkeiten. Math. Ann. 71, 97–115 (1911)
5. Schauder, J.: Der Fixpunktsatz in Funktionalräumen. Studia Math. 2, 171–180 (1930)
6. Tarski, A.: A lattice-theoretical fixpoint theorem and its applications. Pac. J. Math. 5(2), 285–309 (1955)
7. Rudin, W.: Principles of Mathematical Analysis, 3rd edn. McGraw-Hill, New York (1976)
8. Kannan, R.: Some results on fixed points. Bull. Calcutta Math. Soc. 60, 71–76 (1968)
9. Kirk, W.A., Sims, B. (eds.): Handbook of Metric Fixed Point Theory. Springer, Dordrecht (2001)
10. Sperner, E.: Neuer Beweis für die Invarianz der Dimensionszahl und des Gebietes. Abh. Math. Semin. Univ. Hambg. 6, 265–272 (1928)
11. Smart, D.R.: Fixed Point Theorems. Cambridge University Press, Cambridge (1974)
12. Winskel, G.: The Formal Semantics of Programming Languages: An Introduction. MIT Press, Cambridge, MA (1993)
13. Kakutani, S.: A generalization of Brouwer's fixed point theorem. Duke Math. J. 8(3), 457–459 (1941)
14. Nash, J.F.: Equilibrium points in n-person games. Proc. Natl. Acad. Sci. U.S.A. 36(1), 48–49 (1950)
15. Arrow, K.J., Debreu, G.: Existence of an equilibrium for a competitive economy. Econometrica 22(3), 265–290 (1954)
16. Nadler, S.B.: Multi-valued contraction mappings. Pac. J. Math. 30(2), 475–488 (1969)